



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

idius, (3) *Dicrurus modestus atactus*, (4) *Fraseria prosphora*. Two new genera, *Horizocerus* and *Stelgidillas*, are also characterized, and a specimen of the rare Hawk, *Dryotriorchis spectabilis* is reported, the ninth specimen of this species thus far known.—J. A. A.

New Birds from the Bahamas.—Mr. C. J. Maynard, in an 'Appendix to Catalogue of the Birds of the West Indies' (which Catalogue we have not yet seen) has published (Nov. 29, 1899) descriptions of four new species of birds from the Bahamas, namely; (1) *Colinus bahamensis*, from the island of New Providence; (2) *Speotyto bahamensis*, from "New Providence and probably Eleuthera"; (3) *Dendroica bahamensis* ("similar to *Dendroica vigersii*"), from New Providence; (4) *Hæmotopus* (sic) *pratii* (provisional name), from Flemming's Key.—J. A. A.

Kopman on the Bird Fauna of Two Sections of Louisiana.¹—This paper gives a comparison of the representation of 67 species in contiguous but very different portions of southern Louisiana, the fertile alluvial coast district and the pine barrens to the northward. These two areas are separated by the chain of lakes formed by Lake Maurepas, Pontchartrain and Borgne, and mark an abrupt transition from the alluvial fertile district, with its deciduous arboreal vegetation, to the pine districts, or 'pine barrens,' which extend from the eastern border of Louisiana into Mississippi. While scarcely a dozen species are restricted to either of these areas, the relative number of individuals of birds which are common to both varies so greatly as to form a strong contrast in the general ornithological character of the two regions, obviously due to the difference in vegetation and coincident conditions of environment. The birds listed for comparison are mainly the commoner summer residents.—J. A. A.

Faxon and Hoffmann's Birds of Berkshire County, Mass.²—Berkshire County, Massachusetts, differs so much from the rest of the State in altitude and other physical conditions as to form a well-marked region, and one, moreover, until recently ornithologically very imperfectly known. With a general altitude of 1500 to 2000 feet, and with peaks rising from 2400 to 3500 feet, the general character of the fauna and flora is distinctly more boreal than that of that portion of the State to the eastward. For many years ornithologists were left to conjecture as to the birds frequenting the higher parts of 'The Berkshires.' As early as 1884, however, definite

¹ The Bird Fauna of Two Sections. By Henry H. Kopman. The Gulf Fauna and Flora Bulletin, Vol. I, No. 2, pp. 50-57. Dec. 15, 1899.

² The Birds of Berkshire County, Massachusetts. By Walter Faxon and Ralph Hoffmann. Coll. of Berkshire Hist. and Sci. Soc., Vol. III, No. 2, pp. 109-166. Also separate, 8vo, pp. 60. Issued Feb. 23, 1900.